

Safety of SHOCKWAVE Therapy

Laurie Edge-Hughes, BScPT, MAnimSt (Animal Physiotherapy), CAFCI, CCRT
Created for the Animal Rehab Division of the Canadian Physiotherapy Association

BACKGROUND

Shockwave originated in the 1960's and 70's for non-invasive treatment of kidney stones and gallstones (lithotripsy). Human and animal studies in the 1980's incidentally observed osteoblastic response patterns that generated an interest in additional therapeutic uses. Since that time, it has been used successfully for over 20 years to manage a variety of orthopedic conditions (Schmitz et al 2015).

TYPES OF SHOCKWAVE (Van der Worp et al 2013, Cleveland et al 2007, Furia et al 2010, Schmitz et al 2015)

Additional information sources:

- Shockwave therapy BC - http://www.shockwavetherapy.ca/about_eswt.htm - What is [ESWT0](#) Accessed Dec 15, 2014.
- Shockwave therapy - <http://www.shockwavetherapy.eu/> Accessed Dec 15, 2014.
- Shockwave Training – www.shockwavetraining.ca (Calgary, AB, Canada – March 17, 2019.)
- Shockwave theory - <http://www.shockwavetherapy.education/index.php/theory/types-of-shockwave> Accessed March 27, 2019.
- Electrotherapy on the Web: <http://www.electrotherapy.org/modality/shockwave-therapies> - Accessed March 25, 2019.
- Prouza O. 2013. Shockwave therapy in musculoskeletal treatment. http://skanlab.no/wp-content/uploads/2013/10/swt_part_1.pdf (accessed Mar 27, 2019).

There are two types of shockwave therapy. Focused shockwave therapy (FSWT) and radial shockwave therapy (RSWT).

FSWT

A pressure field is generated that converges in the adjustable focus at a selected depth in the body tissues, where the maximal pressure is reached. There are three methods for generation of a focused shockwave: electrohydraulic, electromagnetic and piezoelectric. All of these waves are generated in water (inside of the applicator), which allows for a more natural transference of the waves into the body, with limited reflection.

Electrohydraulic shockwaves are high-energy acoustic waves created by the underwater explosion with high-voltage electrode spark discharge. The waves are then focused with a reflector and targeted at the diseased area. It is a true shock wave at all settings.

Electromagnetic shockwaves are created by an electric current passing through a coil to produce a strong magnetic field. A lens is used to focus the waves, and the focal therapeutic point being defined by the length of the focus lens. The amplitude of the focused waves

increases when the acoustic wave propagates towards the focal point. It is a true shock wave at high settings only.

Piezoelectric shockwaves involve a large number of piezocrystals mounted in a sphere that receives a rapid electrical discharge that induces a pressure pulse in the surrounding water that leads to a shockwave. The arrangement of the crystals causes self-focusing of the waves towards the target centre, which leads to a precise focusing and high-energy within a defined field. It is a true shock wave at high-energy settings only.

RSWT

Radial shock waves are produced by more recently developed pneumatic devices. The term 'radial' refers to the diverging pressure field of the RSWT devices, which reach a maximal pressure at the source of generation (as compared to a focal point away from the applicator). Accelerating a projectile, using compressed air through a tube on the end of which an applicator is placed, generates radial shockwaves. The projectile hits the applicator, which transmits the generated pressure wave into the body. (These waves are NOT generated in water).

INDICATIONS

Shockwave has been shown to be effective for the following select musculoskeletal disorders:

- Calcific tendons & tendinopathies (Kim et al 2014, Verstrelen et al 2014, Gerdesmeyer et al 2003, Cacchio et al 2006, Malliaropoulos et al 2017, Becker et al 2015, Gallagher et al 2012, Wang et al 2002, van der Worp et al 2014, Dedes et al 2018, Cacchio et al 2011, Ilieva et al 2012)
- Plantar fasciitis (Chang et al 2012, Crecco et al 2013, Ilieva 2013, Ibrahim et al 2010)
- Non—Union fractures (Birnbbaum et al 2002, Furia et al 2010)
- Stress fractures (Moretti et al 2009, Rompe et al 2010))
- Osteoarthritis (Zhao et al 2012, Frisbie et al 2009, Kawcak et al 2011, Wang et al 2010, 2011, 2012, Dahlberg et al 2005, Mueller et al 2007, Souza et al 2016)
- Low back pain (Noarnicola et al 2018, Nedelka et al 2014)
- Sacroiliac joint pain (Moon et al 2017)
- Coccydynia (Marwan et al 2017)
- Myofascial trigger points (Ramon et al 2015, Walsh et al 2019)
- Necrosis of the femoral head (Wang CJ et al 2005, Wang L et al 2008)

CONTRAINDICATIONS

- Circulatory disorders where bleeding may be a concern (Desai et al 2017).
- Over malignant tumours. However, cancer itself in a removed area is not a contraindication (Crevenna et al 2019), and newer research is pointing towards shockwave as being an adjunct to cancer care due to its role in permeabilization of mammalian cell membranes (Lopez-Marin et al 2018). However, for now it remains a standard contraindication.

- Presence of infection – (Newer research however, is showing that shockwave may be helpful to reduce infection in chronic wound cases – Zhang et al 2018).
- Shockwave is not generally applied to areas or locations overlying the abdomen or chest (where gas or air is present in the body) (Desai et al 2017, Sistermann & Kathagen 1998).
- Over the abdomen or lumbar spine in pregnant patients (Desai et al 2017).

Manufacturer-Promoted Contraindications without backing or contradiction in the literature

- Metabolic conditions whereby the bone may be fragile.
- Over major blood or nerve vessels too close to a treatment areas.
- Within 4 weeks of a cortisone injection to the area being treated. (The concern is that increased circulation may flush out the injected medication – No studies can be found to validate this contraindication)

Previously thought to be contraindications

- Nerve disorders - Shockwave is now being shown to be harmless to nerve and may improve nerve regeneration (Wu et al 2007, Mense & Hoheisel 2013)
- Over epiphyses in young patients - Newer reviews have stated that it is now considered safe (Lohrer et al 2016)
- Over metal implants – In the case of plated fractures, shockwave is still beneficial, with no adverse events related to the presence of a metal plate. (Wang et al 2001)
- In conjunction with corticosteroid treatment – Topic cortisone in conjunction with shockwave enhanced the effectiveness for plantar fasciitis. (Vahdatpour et al 2018) No studies could be found that compared effectiveness of shockwave therapy in conjunction with a corticosteroid injection.

ADVERSE EVENTS

Redness of the skin, bruising, petechiae, hematoma, and transient discomfort with treatment. (Bannuru et al 2014, Schmitz et al 2015)

CONCLUSIONS from a systematic review on efficacy and safety of Extracorporeal Shockwave Therapy (ESWT) (Schmitz et al 2015)

- **ESWT is effective.**
88.5% of the RCTs (randomized controlled trials) on rESWT (radial) and 81.5% of all RCTs on fESWT (focused) had positive outcomes.
- **ESWT is safe.**
There were no reports of serious adverse events in any of the studies included in this analysis
- **For certain orthopedic conditions, RCTs on ESWT were the predominant type of RCT listed in the PEDro database and/or obtained the highest PEDro scores among all investigated treatment modalities.**
Type of RCT and highest PEDro scores (as compared to all other treatment modalities) were fulfilled for plantar fasciopathy, non-calcific supraspinatus tendinopathy, and calcific tendonitis of the shoulder. RCTs for Achilles tendinopathy and lateral epicondylitis also

ranked high. There were not enough RCTs for ESWT to draw meaningful conclusions regarding greater trochanteric pain syndrome, patellar tendinopathy, knee osteoarthritis, long bone fracture, osteonecrosis of the femoral head, proximal hamstring tendinopathy, long bicipital tenosynovitis, myofascial pain syndrome, myogelosis of the masseter muscle, and spasticity.

- There was **no difference in the ‘quality’ of RCTs on ESWT** in PEDro with positive or negative outcomes.
- **Application of local anesthesia adversely affects outcome of ESWT.**
The molecular mechanisms underlying this phenomenon are not yet fully understood, but substantial evidence points to a central role of the peripheral nervous system in mediating molecular and cellular effects of shock waves applied to the musculoskeletal system. These effects could be blocked by local anesthesia. Thus, it is now generally recommended to apply shock waves without local anesthesia to the musculoskeletal system.
- **Application of insufficient energy adversely affects outcome of ESWT.**
RCTs that showed positive outcomes for rESWT & fESWT for calcifying tendonitis of the shoulder used 2.6x more energy flux density (EFD) than studies that showed a negative outcome. For plantar fasciopathy, positive studies used two times the EFD as negative RCTs. A similar finding was also made when comparing studies for Achilles tendinopathy.
There is no scientific evidence in favour of either rESWT or fESWT with respect to treatment outcome.
It appears that success is more dependent upon sufficient EFD (energy flux density – i.e. power) than with the type of ESWT.
- **The distinction between radial ESWT as ‘low-energy ESWT’ and focused ESWT as ‘high-energy ESWT’ is not correct and should be abandoned.**
Different authors have used different thresholds for categorizing ‘high’ and ‘low’ energy. Because there is no consensus in the literature, this distinction appears arbitrary and should be abandoned.
- **There is no evidence that a certain fESWT technology is superior to other technologies.**
Focused shock waves can be produced by electrohydraulic, electromagnetic, and piezoelectric shock wave generators. The RCTs on fESWT in PEDro do not indicate an advantage of a certain fESWT technology over other technologies.
- **An optimum treatment protocol for ESWT** appears to be three treatment sessions at 1-week intervals, with 2000 impulses per session and the highest EFD that can be applied. This recommendation is based on the average number of treatment sessions and the average interval between treatment sessions among all RCTs on ESWT in PEDro. With respect to the EFD of the impulses (to be as high as possible, i.e. what can be tolerated by the individual patient without application of local anesthesia), this recommendation is based on findings of one study on rESWT for plantar fasciopathy with positive outcome and another study on fESWT for calcifying tendonitis of the shoulder with positive outcome that ‘more is better’. There is not a single RCT on ESWT in PEDro, contradicting this ‘more is better’ recommendation.

SAFETY

Several review papers, meta-analyses, or randomized-controlled trials conclude that shockwave therapy is a safe modality for orthopedic conditions (Schmitz et al 2015), tendinopathies (Dedes et al 2018, Li et al 2017, Cacchio et al 2006), plantar fasciitis (Roerdink et al 2017), and osteoarthritis (Wang TS et al 2020).

References:

1. Bannuru RR, Flavin NE, Vaysbrot E, Harvey W, McAlindon T. High-energy extracorporeal shock-wave therapy for treating chronic calcific tendinitis of the shoulder: a systematic review. *Ann Intern Med.* 2014;160(8):542-549.
2. Becker W, Kowaleski MP, McCarthy RJ, Blake CA. Extracorporeal shockwave therapy for shoulder lameness in dogs. *J Am Anim Hosp Assoc* 51: 15-19, 2015.
3. Birnbaum K et al. Use of extracorporeal shock-wave therapy (ESWT) in the treatment of non-unions. A review of the literature. *Arch Orthop Truma Surg* 2002, 122(6): 324 – 330.
4. Cacchio A, Paoloni M, Barile A, et al. Effectiveness of radial shock-wave therapy for calcific tendinitis of the shoulder: single-blind, randomized clinical study. *Phys Ther* 2006;86:672–82.
5. Cacchio et al. Shockwave therapy for the treatment of chronic proximal hamstring tendinopathy in professional athletes. *Am J Sports Med* 2011, 39(1):146 – 153.
6. Cleveland RO et al. Acoustic field of a ballistic shock wave therapy device. *Ultrasound Med Biol* 2007, 33(8):1327-35.
7. Crevenna R, Mickel M, Keilani M. Extracorporeal shock wave therapy in the supportive care and rehabilitation of cancer patients. *Support Care Cancer.* 2019;27(11):4039-4041.
8. Dahlberg J et al. The evaluation of extracorporeal shockwave therapy in naturally occurring osteoarthritis of the stifle joint in dogs. *Vet Comp Orthop Traumatol* 2005, 18(3): 147 – 152.
9. Dedes V, Stergioulas A, Kipreos G, Dede AM, Mitseas A, Panoutsopoulos GI. Effectiveness and Safety of Shockwave Therapy in Tendinopathies. *Mater Sociomed.* 2018;30(2):131-146.
10. Desai M, Sun Y, Buchholz N, et al. Treatment selection for urolithiasis: percutaneous nephrolithomy, ureteroscopy, shock wave lithotripsy, and active monitoring. *World J Urol.* 2017;35(9):1395-1399.
11. Electrotherapy on the Web: <http://www.electrotherapy.org/modality/shockwave-therapies>- Accessed March 25, 2019.
12. Furia JP et al. Shock wave therapy as a treatment of nonunions, avascular necrosis, and delayed healing of stress fractures. *Foot Ankle Clin N Am* 2010, 15(4): 651 – 662.
13. Frisbie et al. Evaluation of the effect of extracorporeal shock wave treatment on experimentally induced osteoarthritis in middle carpal joints in horses. *Am J Vet Res* 2009, 70(4): 449 – 454.
14. Gallagher A, Cross AR, Sepulveda G. The effect of shock wave therapy on patellar ligament desmitis after tibial plateau leveling osteotomy. *Vet Surg.* 2012 May;41(4):482-5.
15. Gerdesmeyer L et al. Extracorporeal shock wave therapy for the treatment of chronic calcifying tendonitis of the rotator cuff: a randomized controlled trial. *JAMA* 2003;290:2573-80.
16. Ilieva EM et al. Radial shock wave therapy in patients with lateral epicondylitis. *Folia Med*
17. (Plovdiv). 2012 Jul-Sep;54(3):35-41.
18. Ilieva EM. Radial shock wave therapy for plantar fasciitis: a one year follow-up study. *Folia Med (Plovdiv).* 2013;55(1):42-48.
19. Kawcak CE et al. Effects of extracorporeal shock wave therapy and polysulfated glycosaminoglycan treatment on subchondral bone, serum biomarkers, and synovial fluid biomarkers in horses with induced osteoarthritis. *Am J Vet Res* 2011, 72(6): 772 – 779.
20. Kim YS et al. Which method is more effective in treatment of calcific tendinitis in the shoulder? Prospective randomized comparison between ultrasound-guided needling and extracorporeal shock wave therapy. *J Shoulder Elbow Surg.* 2014 Nov;23(11):1640-6.

21. Leeman JJ, Shaw KK, Mison MB, Perry JA, Carr A, Shultz R. Extracorporeal shockwave therapy and therapeutic exercise for supraspinatus and biceps tendinopathies in 29 dogs. *Vet Rec* 179 (15):385, 2016.
22. Li W, Zhang SX, Yang Q, Li BL, Meng QG, Guo ZG. Effect of extracorporeal shock-wave therapy for treating patients with chronic rotator cuff tendonitis. *Medicine (Baltimore)*. 2017;96(35):e7940.
23. Lohrer H, Nauck T, Korakakis V, Malliaropoulos N. Historical ESWT Paradigms Are Overcome: A Narrative Review. *Biomed Res Int*. 2016;2016:3850461.
24. López-Marín LM, Rivera AL, Fernández F, Loske AM. Shock wave-induced permeabilization of mammalian cells. *Phys Life Rev*. 2018;26-27:1-38.
25. Malliaropoulos N, Thompson D, Meke M et al. Individualised radial extracorporeal shock wave therapy (rESWT) for symptomatic calcific shoulder tendinopathy: a retrospective clinical study. *BMC Musculoskeletal Disord*. 2017 Dec 6;18(1):513.
26. Marwan Y, Dahrab B, Esmael A, et al. Extracorporeal shock wave therapy for the treatment of coccydynia: a series of 23 cases. *Eur J Orthop Surg Traumatol*. 2017 Jul;27(5):591-598.
27. Moon YE, Seok H, Kim SH, et al. Extracorporeal shock wave therapy for sacroiliac joint pain: A prospective, randomized, sham-controlled short-term trial. *J Back Musculoskeletal Rehabil*. 2017;30(4):779-784.
28. Moretti B et al. Shock waves in the treatment of stress fractures. *Ultrasound Med Biol* 2009, 35(6): 1042 – 1049.
29. Mueller M1, Bockstahler B, Skalicky M, Mlacnik E, Lorinson D. Effects of radial shockwave therapy on the limb function of dogs with hip osteoarthritis. *Vet Rec*. 2007 Jun 2;160(22):762-5.
30. Nedelka T, Nedelka J, Schlenker J et al. Mechano-transduction effect of shockwaves in the treatment of lumbar facet joint pain: comparative effectiveness evaluation of shockwave therapy, steroid injections and radiofrequency medial branch neurotomy. *Neuro Endocrinol Lett*. 2014;35(5):393-7.
31. Notarnicola A, Maccagnano G, Gallone MF, et al. Extracorporeal shockwave therapy versus exercise program in patients with low back pain: short-term results of a randomised controlled trial. *J Biol Regul Homeost Agents*. 2018 Mar-Apr;32(2):385-389.
32. Prouza O. 2013. Shockwave therapy in musculoskeletal treatment. http://skanlab.no/wp-content/uploads/2013/10/swt_part_1.pdf (accessed Mar 27, 2019).
33. Mense S, Hoheisel U. Shock wave treatment improves nerve regeneration in the rat. *Muscle Nerve*. 2013;47(5):702-710.
34. Roerdink RL, Dietvorst M, van der Zwaard B, van der Worp H, Zwerver J. Complications of extracorporeal shockwave therapy in plantar fasciitis: Systematic review. *Int J Surg*. 2017;46:133-145.
35. Rompe JD, Cacchio A, Furia JP, et al. Low-Energy Extracorporeal Shock Wave Therapy as a Treatment for Medial Tibial Stress Syndrome. *Am J Sports Med* 2010 38: 125.
36. Schmitz C, Császár NB, Milz S, et al. Efficacy and safety of extracorporeal shock wave therapy for orthopedic conditions: a systematic review on studies listed in the PEDro database. *Br Med Bull*. 2015;116(1):115-138.
37. Shockwave therapy BC - http://www.shockwavetherapy.ca/about_eswt.htm - What is ESWT0 Accessed Dec 15, 2014.
38. Shockwave therapy - <http://www.shockwavetherapy.eu/> Accessed Dec 15, 2014.
39. Shockwave Training – www.shockwavetraining.ca (Calgary, AB, Canada – March 17, 2019).
40. Shockwave theory - <http://www.shockwavetherapy.education/index.php/theory/types-of-shockwave> Accessed March 27, 2019.
41. Sistermann R, Katthagen BD. Komplikationen, Nebenwirkungen und Kontraindikationen der Anwendung mittel- und hochenergetischer extrakorporaler Stosswellen im orthopädischen Bereich [Complications, side-effects and contraindications in the use of medium and high-energy extracorporeal shock waves in orthopedics]. *Z Orthop Ihre Grenzgeb*. 1998;136(2):175-181. [Abstract only – article in German]
42. Souza AN, Ferreira MP, Hagen SC, Patrício GC, Matera JM. Radial shock wave therapy in dogs with hip osteoarthritis. *Vet Comp Orthop Traumatol*. 2016;29(2):108-14.
43. Vahdatpour B, Mokhtarian A, Raeissadat SA, Dehghan F, Nasr N, Mazaheri M. Enhancement of the Effectiveness of Extracorporeal Shock Wave Therapy with Topical Corticosteroid in Treatment of Chronic Plantar Fasciitis: A Randomized Control Clinical Trial. *Adv Biomed Res*. 2018;7:62. Published 2018 Apr 24.
44. Van der Worp et al. No difference in effectiveness between focused and radial shockwave therapy for treating patellar tendinopathy: a randomized controlled trial. *Knee Surg Sports Traumatol Arthrosc* 2014, 22(9): 2026 – 2032.

45. Van der Worp H, et al. ESWT for tendinopathy: technology and clinical implications. *Knee Surg Sport Traumatol Arthrosc* 2013, 21: 1451-1458.
46. Verstraelen et al. High-energy Versus Low-energy Extracorporeal Shock Wave Therapy for Calcifying Tendinitis of the Shoulder: Which is Superior? A Meta-analysis. *Clin Orthop Relat Res* 2014, 472:2816–2825
47. Walsh R, Kinsella S, McEvoy J. The effects of dry needling and radial extracorporeal shockwave therapy on latent trigger point sensitivity in the quadriceps: A randomised control pilot study. *J Bodyw Mov Ther.* 2019 Jan;23(1):82-88.
48. Wang CJ, Huang HY, Chen HH, Pai CH, Yang KD. Effect of shock wave therapy on acute fractures of the tibia: a study in a dog model. *Clin Orthop Relat Res.* 2001;(387):112-118.
49. Wang CJ, Huang HY, Pai CH. Shock wave-enhanced neovascularization at the tendon-bone junction: an experiment in dogs. *J Foot Ankle Surg.* 2002 Jan-Feb;41(1):16-22.
50. Wang CJ, Wang FS, Huang CC, et al. Treatment for osteonecrosis of the femoral head: comparison of extracorporeal shock waves with core decompression and bone-grafting. *J Bone Joint Surg Am* 2005;87:2380–7.
51. Wang CJ et al. Extracorporeal shockwave shows regression of osteoarthritis of the knee in rats. *J Surg Res* 2010, 171(2): 601-608.
52. Wang CJ et al. Extracorporeal shockwave shows chondroprotective effects in osteoarthritic rat knee. *Arch Orthop Trauma Surg* 2011, 131(8): 1547 – 1553.
53. Wang CJ et al. Extracorporeal shockwave therapy shows time-dependent chondroprotective effects in osteoarthritis of the knee in rats. *J Surg Res* 2012, 178(1): 196 – 205.
54. Wang CJ. Extracorporeal shockwave therapy in musculoskeletal disorders. *J Orthop Surg Res* 2012, 7:11.
55. Wang L et al. Extracorporeal shock wave therapy in treatment of delayed bone-tendon healing. *Am J Sport Med* 2008, 36(2): 340 – 347.
56. Wang TS, Guo P, Li G, Wang JW. Extracorporeal Shockwave Therapy for Chronic Knee Pain: A Multicenter, Randomized Controlled Trial. *Altern Ther Health Med.* 2020;26(2):34-37.
57. Wu YH, Lun JJ, Chen WS, Chong FC. The electrophysiological and functional effect of shock wave on peripheral nerves. *Conf Proc IEEE Eng Med Biol Soc.* 2007;2007:2369-2372.
58. Zhang L, Fu XB, Chen S, Zhao ZB, Schmitz C, Weng CS. Efficacy and safety of extracorporeal shock wave therapy for acute and chronic soft tissue wounds: A systematic review and meta-analysis. *Int Wound J.* 2018;15(4):590-599.
59. Zelle B et al. Extracorporeal shock wave therapy: Current evidence. *J Orthop Trauma* 2010, 24(3) Suppl 1: S66 – S70.
60. Zhao Z et al. Extracorporeal shock-wave therapy reduces progression of knee osteoarthritis in rabbits by reducing nitric oxide level and chondrocyte apoptosis. *Arch Orthop Trauma Surg* 2012, 132(11): 1547 – 1553.